

CORALLIOPHILINAE (NEOGASTROPODA: MURICIDAE) FROM THE MARQUESAS ISLANDS

MARCO OLIVERIO¹

1 Dipartimento di Biologia Animale e dell'Uomo, "La Sapienza" Rome University, Viale dell'Università 32, I-00185 Roma, Italy.

Abstract Fourteen species of Coralliophilinae (Neogastropoda, Muricidae) have been identified in the material collected by the MUSORSTOM 9 expedition to the Marquesas Islands. This coralliophiline fauna appears severely impoverished, compared to other West Pacific areas. At least half of the species (seven) are members of the shallow water fauna, an unusual pattern for the prevalently deep water coralliophilines, which may indicate an increase with depth of the effects of marginality on benthic faunas. One new species *Coralliophila nukuhiva n. sp.* is here described.

Key words Coralliophilinae, Muricidae, Marquesas, Systematics

INTRODUCTION

The Coralliophilinae, a subfamily of Muricidae with approximately 200-250 described living species, are associated with anthozoan cnidarians, from intertidal habitats down to over 1000 m depth. The Indo-Pacific region hosts the majority of the known species, especially in the core of the region, the area included in the Philippine-Borneo-Papua triangle. Yet the marginal areas in the Pacific are of great biogeographical relevance both for shallow and deep water habitats. The Coralliophilinae, successfully radiated in deep waters, an interesting pattern for a group associated with corals. The present revision, along with a similar one based on materials from the Australes Islands (Oliverio, in prep.) is aimed at providing the faunistic data for a better understanding of the biogeography of these ectobiotic gastropods at the margins of the Indo-West Pacific (see Benton & Spencer [eds], 1995). Previous records of coralliophilines from the Marquesas Islands are to be found in Hombron & Jacquinot (1852), Jardin (1858), Rehder (1969, 1985), Salvat & Rives (1975), Tröndlé (1989), and have been reviewed by Tröndlé & Cose (2005).

MATERIALS AND METHODS

This study is based on the material collected during the expedition MUSORSTOM 9, which included a cruise and a land based workshop. The cruise (led by B. Richer de Forges, IRD, onboard R/V *Alis*) sampled a total of 168 stations by dredging and trawling. Of these, 28 stations

in the waters of seven islands, yielded coralliophiline specimens. The land based workshop (led by R. von Cosel, MNHN) was carried out on Ua Huka Island, sampling a total of 40 stations, of which 6 yielded coralliophiline specimens. For details on the expedition and the complete list of the stations see Richer de Forges *et al.* (1999). Additionally, I have had the possibility to examine the Marquesan samples in the private collections of Jean Tröndlé (La Force: who also took part in the land based workshop at Ua Huka) and Michel Boutet (Tahiti).

A total of 14 species have been identified and are treated in the systematic account. Supraspecific taxonomy in the Coralliophilinae is traditionally based, nearly exclusively, on shell morphology and is currently highly unstable. The first attempts to make a phylogenetic framework for the coralliophiline systematics (Oliverio & Mariottini 2001, Oliverio *et al.* 2002) addressed the need for a deep re-examination by refined morphology and molecular markers. The four genera used herein are thus conceived as in the "traditional" (i.e. Kosuge & Suzuki 1985) taxonomy, yet adopting some slight modifications (e.g. Tsuchiya 2000).

One species is described as new to Science. The protoconchs of 3 species are illustrated with SEM images for the first time. When more than one sample is available for a species, the depth range is given as internal range, which is the range within which the species has certainly been collected (compared to the external range, which is the range where the species may have been collected). Comparative reference is made to Oliverio (in press) for distributional and ecological data, or taxonomical notes.

ABBREVIATIONS

AMNH	America Museum of Natural History, New York, USA.
AMS	The Australian Museum, Sydney.
BMNH	Natural History Museum, London, UK.
IMT	Institute of Malacology, Tokyo
KPM	Kanagawa Prefectural Museum, Yokohama
MNHN	Muséum National d'Histoire Naturelle, Paris, France.
SAM	South Africa Museum, Cape Town
SDSNH	San Diego Society of Natural History, California, USA
sh	empty shell(s).
spm	live collected specimen(s).
SSSM	Sea Shell Specimen Museum, Shinjuku, Tokyo
TA	Toba Aquarium, Toba
TwM	Taiwan Museum, Taiwan.
USNM	Museum of Natural History, Smithsonian Institution, Washington DC, USA.
ZMB	Museum of Natural History, Humboldt University, Berlin
ZMUC	Zoological Museum University of Copenhagen

SYSTEMATICS

Superfamily Muricoidea Rafinesque, 1815
 Family Muricidae Rafinesque, 1815
 Subfamily Coralliophilinae Chenu, 1859

Genus *Coralliophila* H. & A. Adams, 1853

Coralliophila H. & A. Adams, 1853: 135. Type species: *Fusus neritoideus* Lamarck, 1816 (= *Purpura violacea* Kiener, 1836), by subsequent designation (Iredale, 1912).

Coralliobia H. & A. Adams, 1853: 138. Type species (by monotypy): *Concholepas (Coralliobia) fimbriata* A. Adams 1854; Recent, Indo-West Pacific.

Galeropsis Hupé, 1860: 125. Type species (by monotypy): *Galeropsis lavenayanus* Hupé, 1860; Lower Miocene, France.

Pseudomurex Monterosato, 1872: 15. Type species (by original designation): *Murex meyendorffii* Calcara, 1845; Recent, Northeastern Atlantic.

Quoyula Iredale, 1912: 221. Type species (by original designation): *Purpura monodonta* Blainville,

1832; Recent, Indo-Pacific.

Reliquiaecava Massin, 1987: 81. Type species (by original designation): *Leptoconchus robilliardi* Liénard, 1870. Recent, Indo-Pacific.

Rhombothais Woolacot, 1954: 38. Type species (by original designation): *Rhombothais arbutum* Woolacot, 1954 Recent, Australia.

Remarks Since *Coralliophila* as traditionally conceived is clearly polyphyletic (Oliverio & Mariottini 2001, and new unpublished data), the genus is here used in a wide sense (*Coralliophila sensu lato*). Also the list of synonymies for *Coralliophila* is deemed to be revised in a phylogenetic framework. I have included in *Coralliophila* (s.l.) also *Rapana nodosa* A. Adams, 1854, which is traditionally included in *Mipus*, following Tsuchiya (2000) and particularly Oliverio (in press).

Coralliophila violacea (Kiener, 1836)

"*Murex neritoideus*" Chemnitz, 1788: 280, pl. 165, figs 1577, 1578 (non binominal).

Murex neritoideus Gmelin, 1791: 3559 (non Linnaeus, 1767).

Fusus neritoideus Lamarck, 1816: pl. 435, figs 2a, b.

Purpura violacea Kiener, 1836: 77, pl. 19, fig. 57.

Type material *Murex neritoideus* Gmelin, Chemnitz's (1788) figured specimen (not found in Copenhagen nor in S. Petersburg).

Type locality *Purpura violacea* Kiener [*Murex neritoideus* Gmelin]: Nicobar Islands, "an den Stranden der Nicobarischen Enlande" (Chemnitz 1788: 280).

Material examined — MUSORSTOM 9: Ua Huka Is.: stn 19, Hane Bay, 8°56.65'S, 139°42.40'W, 0-3 m, 2 spm; Nuku Hiva Is., intertidal 4 sh. (coll. Jean Tröndlé, La Force and Michel Boutet, Tahiti).

Distribution The entire Indo-Pacific. From the Red Sea and the western Indian Ocean to the tropical eastern Pacific, Panamic province to Ecuador, subtidal associated with *Porites* (e.g.: Bosch *et al.* 1995; Tsuchiya 2000; Keen 1971; Cernohorsky 1972). From New Caledonia, alive in 4-25 m (Oliverio in press). Marquesas, alive in 0-3 m.

Remarks This is the type species of *Coralliophila*. The nomenclatural history of *Coralliophila violacea* was summarized by Cernohorsky (1985) and Oliverio (in press). It has been reported from Marquesas by Jardin (1858: 188) and by Rehder (1969: 31).

Coralliophila bulbiformis (Conrad, 1837)

Purpura bulbiformis Conrad, 1837: 266, pl. 20, fig. 23.

Purpura gibbosa Reeve, 1846: sp. 78.

Coralliophila cantrainei Montrouzier in Souverbie, 1861: 282, pl. 11, fig. 11.

Coralliophila elaborata H. & A. Adams, 1863: 433.

Coralliophila nivea Oliver 1915: 536 (non A. Adams, 1853).

Coralliophila erosa Kay 1979: 255, fig. 90B (non Röding, 1798).

Type material *Purpura bulbiformis*: not found. — *Purpura gibbosa*: 3 possible syntypes BMNH 1984102, none of which matches the original illustration (K.M. Way pers. comm.), one figured by Kosuge & Suzuki (1985: pl. 47, fig. 2). — *Coralliophila cantrainei*: Kosuge & Suzuki (1985: pl. 47, fig. 1) figured an alleged "syntype" from MNHN; actually there are 4 specimens "ex. d'auteur" (determined by the author - Montrouzier - yet not type material) in MNHN, the holotype should be in the Musée de Bordeaux. — *Coralliophila elaborata*: lectotype ("type", Kosuge & Suzuki 1985, pl. 37, fig. 8) BMNH 1982057/1 and 2 paralectotypes BMNH 1982057/2-3.

Type locality *Purpura bulbiformis*: Kauai [Atooi], Hawaiian Islands. — *Purpura gibbosa*: no locality reported. — *Coralliophila cantrainei*: Art Island, New Caledonia. — *Coralliophila elaborata*: Hawaii.

Material examined — Nuku Hiva Is., intertidal 2 sh. (coll. Jean Tröndlé, La Force); Fatu Hiva Is., 1 sh. (coll. Michel Boutet, Tahiti).

Distribution Throughout the Indo-West Pacific, intertidal with hermatypic corals (Cernohorsky 1978; Brook 1999: 453). From New Caledonia, alive in 4-32 m (Oliverio in press). Marquesas, shell only, intertidal.

Remarks It was reported from Marquesas by Salvat & Rives [under *C. costularis* (Lamarck, 1816)], by Rehder (1985: 98) and by Tröndlé (1989: 19).

Coralliophila abnormis (Smith, 1878)

Figures 1-3, 30

Fusus abnormis Smith, 1878: 811-812, pl. 50, fig. 10.

Latiaxis chiangi Lan, 1982: 23, fig. 1.

Coralliophila armeniaca D'Attilio & Myers, 1984: 89, figs 21-24.


Type material *Fusus abnormis*: holotype, BMNH 1878.6.10.17, figured by Kosuge & Suzuki (1985, pl. 39, fig. 1) and Higo et al. (2001: 66, fig. G2403). — *Latiaxis chiangi*: holotype, TwM MT8218. — *Coralliophila armeniaca*: holotype, SDSHN 79507 and 14 paratypes (SDSHN 79504a-j, AMNH 213803a-b, USNM 819634a-b).

Type locality *Fusus abnormis*: Andaman Islands, off Port Blair. — *Latiaxis chiangi*: Bohol Island, Philippines. — *Coralliophila armeniaca*: Bohol Straits, 75 m depth.

Material examined MUSORSTOM 9: Ua Pou Is., stn DW 1144, 9°19.3'S, 140°03.8'W, 85-95 m, 2 sh; Eiao Is., stn DW 1154, 7°58.5'S, 140°43.7'W, 102 m, 1 sh; Nuku Hiva Is., stn DW 1170, 8°45.1'S, 140°13.1'W, 104-109 m, 9 sh; stn DR 1181, 8°45.5'S, 140°03.2'W, 102-130 m, 1 spm 3 sh; stn DR 1182, 8°45.6'S, 140°03.9'W, 90-120 m, 3 sh; stn DR 1305, 8°54.1'S, 140°14.5'W, 90-155 m, 2 sh; Hiva Oa Is., stn DW 1208, 9°48.9'S, 139°09.5'W, 117-117 m, 1 sh; stn DW 1224, 9°44.6'S, 138°51.1'W, 115-120 m, 1 sh; Fatu Hiva Is., stn DW 1242, 10°28.1'S, 138°41.1'W, 119-122 m, 1 juv sh; Ua Huka Is., stn DR 1292, 8°54.1'S, 139°37.8'W, 95-100 m, 1 sh.

Distribution Indian Ocean, Somalia (Kosuge 1993) to the Andaman Islands (Smith 1878). Pacific Ocean, from Kii Peninsula to the Philippines (Tsuchiya 2000). From New Caledonia, alive in 15-25 m (Oliverio in press). Marquesas, one alive in 102-130 m, empty shells in 95-119 m..

Remarks Protoconch of 3.8 whorls (Fig. 30), 700 µm high and 650 µm wide at the base. Protoconch I of 0.9 whorls, sculptured with pustules over the entire



Figures 1-10 Shells of Coralliophilinae. **1-3** *Coralliophila abnormis* (Smith, 1878): **1** Nuku Hiva Is., stn DW 1170, 104-109 m [15.2×8.8mm]; **2-3** - stn DR 1181, 102-130 m [14.4×7.5 mm]. **4** *Coralliophila clathrata* (A. Adams, 1854), Ua Huka Is.: stn 20, South coast, Hinipahué Bay, 8°56.20'S, 139°32.90'W, intertidal, [11.5×8.1 mm]. **5** *Coralliophila monodonta* (Blainville, 1832), Ua Huka Is., stn 23, Haamamao Bay, intertidal [20.5×14.7 mm]. **6, 7** *Coralliophila curta* G.B. Sowerby III, 1894: Hiva Oa Is., stn DW 1204, 60-62 m [7.6×4.8 mm]. **8-10** *Coralliophila nukuhiva* n. sp.: **8, 9**, Holotype stn CP 1176, Nuku Hiva Is., 260 m. (MNHN 20896), [20.8×15 mm]. **10** New Caledonia, BATHUS 4 stn CP 928 [24.0×15.5 mm].


surface (partly abraded in the figured specimen). Protoconch-II of 2.9 whorls, with two major spiral keels, and a minor subsutural one, each sculptured by a series of threads (prosocline on the minor subsutural keel, opisthocline on the major keels). The threads tend to fuse, giving rise to collabral riblets. This morphology may be considered as grossly concordant with the protoconch of *Coralliophila armeniaca*, described by D'Attilio & Myers (1984: 89, fig. 23) as multispiral (3.5 whorls) with two poorly nodulose spiral cords crossed by axial ridges. *Coralliophila abnormis* is normally found throughout its range in water shallower than about 100 m and is thus considered a shallow water species (Oliverio in press). All present specimens were collected in slightly deeper waters at Marquesas where passive transport cannot be excluded, even for the specimen collected alive.

***Coralliophila clathrata* (A. Adams, 1854)**
Figure 4

Rapana (Rhizochilus) clathrata A. Adams in H. & A. Adams 1853-58 [1854]: 97.
Coralliobia sugimotonis Kuroda, 1931: 316, pl. 1, figs 5-7.

Type material *Rapana clathrata*: lectotype (Robertson 1980: 8, figs 8, 9) BMNH 1882.12.6.73, figured by Kosuge & Suzuki (1985: pl. 46, fig. 4). — *Coralliobia sugimotonis*: not found.

Type locality *Rapana clathrata*: Philippines (Robertson 1980). — *Coralliobia sugimotonis*: Okinoshima, Tosa Bay, Japan.



Figures 11-20 Shells of Coralliophilinae. **11, 12** *Coralliophila mitraeformata* Kosuge, 1985: Fatu Hiva Is., stn DR 1247, 1150-1250 m [Fig. 9 4×4.6 mm]. **13-15** *Coralliophila nodosa* (A. Adams, 1854): Nuku Hiva stn DW 1170, 104-109 m [Figs 13, 14: 16.9×12.4 mm,] [Fig. 15: 12×8.6 mm]. **16-20** *Rhizochilus cf. antipathum* Steenstrup, 1850: Ua Pou Is., stn CP 1159, 145 m [Fig. 17 7.5×4.4 mm]

Material examined MUSORSTOM 9: Ua Huka Is.: stn 20, South coast, Hinipahué Bay, 8°56.20'S, 139°32.90'W, intertidal, 2 spm, 1 sh; stn 19, Hane Bay, 8°56.65'S, 139°42.40'W, 0-3 m, 1 sh; stn 23, Haamamao Bay, 8°55.90'S, 139°31.45'W, intertidal, 1 sh.

Ua Pou Is., intertidal, 1 sh (coll. Jean Tröndlé, La Force); Nuku Hiva Is., intertidal, 2 sh. (coll. Jean Tröndlé, La Force), 2 sh (coll. Michel Boutet, Tahiti); Fatu Hiva Is. 1 spm, 1 sh (coll. Michel Boutet, Tahiti : ex. coll. G. Richard).

Distribution Indo-Pacific. Indian Ocean, from Natal to Sri Lanka and Maldives Islands (Barnard 1969: 637, fig. 17c, as *Coralliophila* sp.; Robertson 1980). Pacific Ocean, from Kii Peninsula (Japan) southward to the Great Barrier Reef, through the tropical Pacific, eastward to the Marquesas Islands (Robertson 1980; Tsuchiya 2000). New

Caledonia, shells only in 4-20 m (Oliverio in press). Marquesas, living intertidal, empty shells down to 3 m.

Remarks It is known to be associated with, and to feed on, the zoanthids *Palythoa* spp. (Robertson 1980). It was reported from Marquesas by Tröndlé (1989).

Coralliophila monodonta (Blainville, 1832)


Figure 5

Purpura monodonta Blainville, 1832: 241.

Purpura monodonta Quoy & Gaimard, 1833: 561, pl. 37[1834], figs 9-11.

Purpura madreporarum G.B. Sowerby I in J. Sowerby & G.B. Sowerby I, 1834: pl. 237, fig. 12.

Quoyula madreparium [sic] - Oliver 1915: 537.



Figures 21-29 Shells of Coralliophilinae. **21, 22** *Babelomurex japonicus* (Dunker, 1882), Nuku Hiva Is., stn CP 1176, 260-260 m [Fig. 21: 18×14.9 mm]. **23-27** *Babelomurex takahashii* (Kosuge, 1979). 23, 24 Nuku Hiva Is., stn DW 1170, 104-109 m [Fig. 23: 19.5×13.2 mm; Fig. 24: 31×21.5 mm]. 25, Hiva Oa Is., stn DW 1224, 115-120 m [SEM 9.8×5.7 mm]. 26, 27, Holotype of *Babelomurex cookae* USNM 859322, Hawaii Islands [39.0×24.7 mm]. **28, 29** *Leptoconchus lamarckii* Deshayes, 1863, Ua Pou stn DW 1143, 18-55 m [Fig. 24.1×11.1 mm].

Coralliophila madreporara [sic] - Kosuge & Suzuki 1985: 35, pl. 46, fig. 7. — Bosch *et al.* 1995: 125, fig. 501.

Galeropsis madreporarus [sic] - Lozouet & Renard 1998: 181, figs 7/4, 5.

Coralliophila madreporaria [sic] - Tsuchiya 2000: 419, pl. 208, fig. 291.

Type material *Purpura monodonta* Blainville: lectotype (Oliverio in press) and 1 paralectotype MNHN. — *Purpura madreporarum*: 3 possible syntypes BMNH 1985104.

Type locality *Purpura monodonta* Blainville: Tonga-Tabou. — *Purpura madreporarum* Sowerby G.B. I: unknown.


Material examined MUSORSTOM 9: Nuku Hiva Is., stn CP 1177, 8°45.1'S, 140°14.1'W, 108-112 m, 1

spm; Hiva Oa Is., stn DW 1206, 9°51.4'S, 139°09.1'W, 352-358 m, 1 juv sh; Fatu Hiva Is., stn DR 1244, 10°28.4'S, 138°42.1'W, 1015-1020 m, 1 sh.

Ua Huka Is.: stn 1, Hane Bay, Hinipahué Bay, 8°55.6'S, 139°32.1'W, intertidal, 1 sh; stn 23, Haamamao Bay, 8°55.90'S, 139°31.45'W, intertidal, 1 sh; stn 24bis, Haahue Bay, 8°53.60'S, 139°37.0'W, 20-33 m, 1 sh.

Rimatala Is. intertidal, 1 sh (coll. Jean Tröndlé, La Force); Ua Pou Is., intertidal, 1 sh (coll. Jean Tröndlé, La Force); Nuku Hiva Is., intertidal, 1 sh (coll. Jean Tröndlé, La Force).

Distribution Indo-Pacific, from the Red Sea and the eastern Africa to Indonesia, throughout the tropical Pacific (e.g.: Bosch *et al.* 1995; Cernohorsky 1972), including Hawaii (Kay 1979) and Easter Island (Rehder 1980), and extending in the eastern Pacific to the Galapagos Islands



Figures 30-32 Protoconch of Coralliophilinae. **30** *Coralliophila abnormis* (Smith, 1878): Nuku Hiva Is., stn DW 1170, 104-109 m. **31** *Coralliophila nodosa* (A. Adams, 1854): Nuku Hiva stn DW 1170, 104-109 m. **32** *Babelomurex takahashii* (Kosuge, 1979), Hiva Oa Is., stn DW 1224, 115-120 m. Scale bars 200 µm.

(Finet 1994) and the American west coast (Keen 1971). From New Caledonia, alive in 4-40 m (Oliverio in press). Marquesas, empty shells intertidal to 33 m, with deep samples (empty shells 112-1015 !) likely to be the result of drifting down the slope.

Remarks The question of the valid name for this species is dealt with by Oliverio (in press), who established that the binomen *Purpura monodonta* Blainville was dated to 1832 and was thus the first available. Recorded from Marquesas by Hombron & Jacquinot (1852), Jardin (1858) and Rehder (1969), all under *Purpura madreporarum*, and by Tröndlé & Cosel (2005).

***Coralliophila curta* G.B. Sowerby III, 1894**
Figures 6, 7

Coralliophila curta G.B. Sowerby III, 1894: 42, pl. 4, fig. 4.

Type material Holotype BMNH 1890.9.25.23, figured by Smriglio & Mariottini (2000: 13, figs 9, 10).

Type locality Mauritius.

Material examined MUSORSTOM 9: Hiva Oa Is., stn DW 1204, 9°52.6'S, 139°03.2'W, 60-62 m, 1 sh.

Nuku Hiva Is., intertidal, 1 sh (coll. Michel Boutet, Tahiti).

Distribution Indo-West Pacific, from Mauritius to southern Japan, the Philippine and New

Caledonia (Kosuge & Suzuki 1985; Tsuchiya 2000; Oliverio in press), intertidal to 30 m. Marquesas, empty shells intertidal to 60 m.

Remarks The protoconch of 2.6 conical whorls (Oliverio in press) indicates a planktotrophic development. This record, the easternmost for the species, extends the range to the entire western Pacific.

***Coralliophila nukuhiva* n. sp.**
Figs 8-10

Holotype 1 spm (Figs 8, 9), from stn CP 1176, Nuku Hiva Is., Marquesas, 8°44.8'S, 140°14.5'W, 260 m, MNHN 20896.

Paratypes Ua Huka Is., stn DW 1288, 8°53.9'S, 139°38.0'W, 200-220 m, 1 spm, MNHN 20897; stn CP 1300, 8°49.9'S, 140°17.4'W, 416-430 m, 1 spm, MNHN 20898; Hiva Oa Is., stn DR 1198, 9°50.0'S, 139°09.4'W, 290-320 m, 1 spm, MNHN 20899; stn DW 1201, 9°50.6'S, 139°09.2'W, 275-300 m, 3 spm, 1 sh, MNHN 20900; Hatutaa Is., stn DW 1287, 7°54.5'S, 140°40.2'W, 163-245 m, 1 spm, MNHN 20901.

Type locality Nuku Hiva Is., Marquesas, 8°44.8'S, 140°14.5'W, 260-260 m, [MUSORSTOM 9: stn CP 1176].

Material examined Marquesas MUSORSTOM 9: Ua Huka Is., stn DW 1288, 8°53.9'S, 139°38.0'W, 200-220 m, 1 sh. North of New Caledonia BATHUS 4: stn DW 927, Grand Passage, 18°56'S, 163°22'E, 444-452 m, 1 sh; stn CP 928, 18°55'S,

163°24'E, 420-452 m, 1 sh (Fig. 10). **Norfolk Ridge** BATHUS 3: stn CP 811, Banc Jumeau Ouest, 23°41'S, 168°15'E, 383-408 m, 1 sh.

Measurements Size of the type specimens (height × width, mm). Holotype: 20.8×15. Paratypes: 22.1×14.4, 21.2×13.3, 12×8.1, 14×8.6, 11.2×7.1, 12.3×7.5, 13.9×9.1, 12×7.6. Average height/width ratio 1.5 (std 0.08).

Description Shell of medium size for genus, height 20.8 mm and width 15 mm, thick, rhomboidal in shape, with angled shoulder.

Protoconch missing in the holotype. *Teleoconch* of 4.2+ whorls (tip missing in the holotype). Spire relatively high, conical, with flat sides, suture very incised. Aperture more than half of total height, last whorl over $\frac{3}{4}$ of total height. Spire strongly curved at the shoulder, base constricted. Aperture long, narrowly oval. Outer lip externally crenulated. Inner lip straight, callous. Siphonal canal moderately long, open; umbilical area moderately wide with imbricate fasciole, umbilicus narrowly open. Sculpture on the teleoconch whorls consisting of closely set spiral cords with minute, densely packed imbricate scaly spines. One cord on the first whorl, then additional cords appearing, gradually increasing in size. Five major cords and two minor cordlets above the aperture and a total of 23 cords on the last whorl, 17 major and 6 minor, irregularly alternating. Axial sculpture of 8 broad, slightly prosocline ribs per whorl. Ground colour ivory white, inside the aperture bright white.

Animal and operculum unknown.

Derivation of name The epithet *nukuhiva* refers to the island (Nuku Hiva Is.) where the holotype was collected. Used as a noun in apposition.

Distribution *C. nukuhiva* n. sp. is known from the Marquesas, alive in 220-416 m, and New Caledonia and Norfolk Ridge, empty shells in 408-444 m.

Remarks The protoconch was totally missing in all specimens except for a remnant in one paratype. This much eroded whorl of the protoconch had traces of two keels and sinusigera lip, indicating a multispiral type and thus a planktotrophic development. *C. nukuhiva* is similar to *C. nivea* (A. Adams, 1853) from which it differs in its

coarser spiral sculpture of more closely set cords, its less slender spire, with a more pronounced shoulder and incised suture. From *Coralliophila ovoidea* (Kosuge, 1985) it differs in its less elongated spire, proportionally higher aperture and more stepped shoulder. It is also similar to *C. fritschii* (Martens, 1874), from South Africa, in the pattern of teleoconch sculpture but differs in its less slender outline, in having coarser sculpture and a less constricted base. From *C. roseocephala* Kosuge, 1986, it differs in its less slender spire with more convex whorls.

Coralliophila mitraeformata Kosuge, 1985

Figs 11, 12

Coralliophila mitraeformata Kosuge, 1985: 56, pl. 21, figs 1-2.

Type material Holotype IMT 85-21, figured by Higo et al. (2001: 66, G2401).

Type locality Mactan, Cebu, Philippines.

Material examined MUSORSTOM 9: Fatu Hiva Is., stn DR 1247, 10°34.0'S, 138°41.6'W, 1150-1250 m, 1 sh.

Distribution Pacific Ocean, from off Danjo Islands, Western Ki sh to the Philippines (Higo et al. 1999: 217, Tsuchiya 2000). From New Caledonia, alive in 245-400 m (Oliverio in press). Marquesas, one empty shell in 1150-1250 m.

Remarks The protoconch, which was partly eroded and of 2.5+ conical whorls, indicates a planktotrophic larval development. This species is very similar to *C. solutistoma* Kuroda & Shikama, 1966, the main difference being the higher value of the ratio "total height/apertural height" (h/a). Specimens with a ratio of 1.7 or less exist and these can hardly be separated from slender specimens of *C. solutistoma*. Specimens with a ratio greater than 1.7 are easily identified as *C. mitraeformata*. This record significantly extends the range eastward in the Pacific.

Coralliophila nodosa (A. Adams, 1854)

Fig. 13-15, 31

Rapana (Latiaxis) nodosa A. Adams, 1854: 98.

Type material Two syntypes, BMNH 1984114, figured by Kosuge & Suzuki (1985: pl. 25, fig. 8, pl. 46, fig. 10).

Type locality Philippines.

Material examined MUSORSTOM 9: Nuku Hiva stn DW 1170, 8°45.1'S, 140°13.1'W, 104-109 m, 1 spm, 2 sh.

Distribution Western Pacific, Philippines to Hawaii (Kay 1979). New Caledonia, empty shells in 230-266 m (Oliverio in press). Marquesas alive in 104-109 m.

Remarks A single, partly eroded (last 2.6 whorls remaining) measured 720 µm wide at the base and 790+ µm high (Fig. 31). Protoconch I is missing. Protoconch II has two major spiral keels covered by nodulose tubercles and a series of subsutural tubercles visible. A sinusigera lip at the protoconch-teleoconch boundary is present.

The referral of the present shells to Adam's taxon is not definitive. The general outline is similar to that of *Coralliophila crebrilamellosa*, the main difference being the less scabrous spiral sculpture. The specimens listed here seem constant in their characters. *Rhombothais arbutum* Woolacot, 1954, from Australia, and *Latiaxis rosaceus* Smith, 1903, from South Africa, are similar. All are commonly placed in *Mipus*, and possibly represent a complex with an Indo-West Pacific distribution.

***Coralliophila fimbriata* (A. Adams, 1854)**
Figs 47, 163

Concholepas (Coralliobia) fimbriata A. Adams, 1854: 93.

Coralliobia cancellata Pease, 1861: 399.

Coralliobia sculptilis Pease, 1865: 513.

Coralliobia smithi Yen, 1942: 226.

Coralliobia densicostata Shikama, 1963

Type material *Concholepas fimbriata*: syntype BMNH 1985103, figured by Kosuge & Suzuki (1985: pl. 48, fig. 8). — *Coralliobia sculptilis*: not found (Kay 1965: 86). — *Coralliobia cancellata*: not found (Kay 1965: 85). — *Coralliobia smithi*: holotype BMNH 94.4.5.17, figured by Kosuge & Suzuki (1985: pl. 45, fig. 5). — *Coralliobia densicostata*: syntype KPM NG0120043, figured

by Kosuge & Suzuki (1985: pl. 45, fig. 6) and by Higo *et al.* (2001: 66, G2411).

Type locality *Concholepas fimbriata*: Cagayan, Mindanao, Philippines. — *Coralliobia sculptilis*: Hawaiian Islands [Sandwich Islands]. — *Coralliobia cancellata*: Hawaiian Islands [Sandwich Islands]. — *Coralliobia smithi*: Macclesfield Bank, China Sea. — *Coralliobia densicostata*: Nase, Amami-Oshima, Japan.

Material examined MUSORSTOM 9: Hiva Oa Is., stn DW 1204, 9°52.6'S, 139°03.2'W, 60-62 m, 1 sh.

Distribution Tropical and subtropical Indo-West Pacific (Kay 1979; Kosuge & Suzuki 1985). New Caledonia, alive in 90-305 m, empty shells in 12-447 m. Present material, alive in 90-367 m, empty shells in 5-487 m.

Remarks *Coralliophila fimbriata* is the type species of *Coralliobia* H. & A. Adams, 1853, but I prefer to keep it in *Coralliophila* [s.l.] until the systematics of coralliophilines is clarified and relationships with other nominal genera (e.g. *Emozamia*) have been reviewed. *Coralliophila fimbriata* is associated with hermatypic scleractinians, yet it has frequently been collected alive at depths well over 100 m.

Genus *Rhizochilus* Steenstrup, 1850

Rhizochilus Steenstrup, 1850: 75. Type species (by monotypy): *Rhizochilus antipathum* Steenstrup, 1850; Recent, Indo-Pacific.

***Rhizochilus* cf. *antipathum* Steenstrup, 1850**
Figs 16-20

Rhizochilus antipathum Steenstrup, 1850: 75.

Rhizochilus teramachii Kuroda, 1953: 118, 125-126, text figs 1, 2.

Type material *Rhizochilus antipathum*: 7 syntypes ZMUC, one figured by Kosuge & Suzuki (1985: pl. 50, fig. 6). — *Rhizochilus teramachii*: holotype in TA 1921, figured by Kosuge & Suzuki (1985: pl. 45, fig. 12) and Higo *et al.* (2001: 66, G2412s).

Type locality Rhizochilus antipathum: No locality is given either with the type material (Ole Tendal, pers. comm.) or in the original description.— *Rhizochilus teramachii:* East of Ashizuri-Misaki, Japan, 80-90 fms [146-165 m].

Material examined MUSORSTOM 9: Ua Pou Is., stn CP 1159, 7°58.3'S, 140°43.7'W, 145 m, 1 spm, 1 juv spm; Eiao Is., stn CP 1271, 7°53.6'S, 140°42.2'W, 600 m, 2 spm (on coral branch); stn DW 1274, 7°54.6'S, 140°40.1'W, 100-120 m, 2 spm (on coral branch).

Distribution Indo-Pacific, Indian Ocean (Réunion Is., MO, unpublished) and western Pacific, from Japan, (Tsuchiya 2000), to Hawaii (Gage 1962; Kay 1979). New Caledonia, shells only in 25-400 m. Marquesas, alive in 120-600 m.

Remarks The multispiral protoconch of 3.2 conical whorls is indicative of a planktotrophic development. This species was recorded for the Marquesas by Tröndlé (1989: 25, referring to material from the PELE expedition based on a pers. comm. by H.A. Rehder). All recorded associations refer to antipatharians (e.g. "on *Antipathes japonica* Brook": Higo *et al.* 1999: 217, G2412; on *Antipathes grandis*: Gage 1962). Beyond the typical deformation in the adults, there is some variability in shell features including colour and sculpture. I cannot exclude the possibility that more than one species is involved, with different black coral hosts.

Genus *Babelomurex* Coen, 1922

Babelomurex Coen, 1922: 68. Type species (by original designation): *Fusus babelis* Requier, 1848; Recent, north-eastern Atlantic and Mediterranean.

Langfordia Dall, 1924: 89. Type species (by original designation): *Murex cuspidifera* Dall, 1924. *Tolema* Iredale, 1929: 186. Type species (ICZN, 1970): "*Tolema australis* Laseron, 1955" (incorrect name used = *Rapana lischkeanus* Dunker, 1882).

Remarks The species of *Babelomurex* have been classified in the past into several subgenera, based on shell features (e.g. Kosuge & Suzuki 1985; Higo *et al.* 1999, 2000). I have preferred a more conservative position, pending the establishment of a reli-

able phylogenetic scheme for these putative taxa.

Babelomurex japonicus (Dunker, 1882)

Figs 21-22

Rapana japonica Dunker, 1882: 43, pl. 13, figs 24-25.

Latiaxis sallei Jousseaume, 1884: 186-187, pl. 10, fig. 3.

Type material *Rapana japonica*: holotype ZMB 103.966, figured by Kosuge & Suzuki 1985: pl. 27, fig. 1. — *Latiaxis sallei*: holotype MNHN, figured by Kosuge & Suzuki 1985: pl. 27, fig. 2.

Type locality *Rapana japonica*: Seto Inland Sea, Japan. — *Latiaxis sallei*: Japan.

Material examined MUSORSTOM 9: Nuku Hiva Is., stn CP 1176, 8°44.8'S, 140°14.5'W, 260-260 m, 2 sh.

Distribution West Pacific, from Japan to the Philippines and Australia, and through the Pacific to Hawaii (Cernohorsky 1978; Kay 1979; Tsuchiya 2000). From New Caledonia, alive in 250-253 m (Oliverio in press). Marquesas, empty shells in 260 m.

Remarks Diagnosis of the nominal taxa included in the *Babelomurex japonicus* group is traditionally based on the direction of the spines, the scabrousness of the spiral sculpture, the slenderness of the shell and the presence or absence of a scabrous ridge on the shoulder. There seems to be some variation throughout the range in teleoconch features and also in dimensional parameters of the protoconchs. The two shells of the present material are, as seems common in the SW Pacific *B. japonicus*, smaller than those from the Philippines. Shells of *B. japonicus* with an intact apex that I have examined from throughout the range always had a multispiral protoconch of 3.2-3.3 conical whorls indicating a planktotrophic larval development.

Babelomurex takahashii (Kosuge, 1979)

Figs 23-27, 32

Latiaxis (Echinolatiaxis) takahashii Kosuge, 1979: 4, pl. 3, figs 4-5.

Babelomurex cookae Kosuge, 1988: 99 pl. 39 figs 1-4

Type material *Latiaxis takahashii*: Holotype IMT 79-7, figured by Kosuge & Suzuki (1985: pl. 16, fig. 13, pl. 33, fig. 4). — *Babelomurex cookae*: Holotype USNM 859322

Type locality *Latiaxis takahashii*: “Of Mactan Id., Cebu, Philippines, at about 100 m deep” (Kosuge, 1979: 5). — *Babelomurex cookae*: “French Frigate Shoals, 525 miles N. E. of Ohau Island, Hawaii Islands at the depth of about 75 to 175 fathoms” [136-318 m] (Kosuge, 1988: 100).

Material examined MUSORSTOM 9: Nuku Hiva Is., stn DW 1170, 8°45.1'S, 140°13.1'W, 104-109 m, 15 sh; stn CP 1177, 8°45.1'S, 140°14.1'W, 108-112 m, 1 sh; stn DR 1182, 8°45.6'S, 140°03.9'W, 90-120 m, 1 sh; stn DR 1183, 8°45.5'S, 140°03.8'W, 86-120 m, 1 sh; Hiva Oa Is., stn DW 1224, 9°44.6'S, 138°51.1'W, 115-120 m, 1 sh; Ua Pou Is., stn CP 1265, 9°20.4'S, 140°07.3'W, 90-92 m, 1 spm, 1 sh; Nuku Hiva Is., stn DR 1298, 8°49.1'S, 140°17.1'W, 305 m, 1 sh; stn DR 1305, 8°54.1'S, 140°14.5'W, 90-155 m, 1 sh.

Distribution Pacific Ocean, from southern Japan (Ōsumi Islands) to the Philippines, 50-120 m (Higo *et al.* 1999: 220), eastward to the Hawaii in 136-318 m (as *B. cookae*). Marquesas, alive in 90-92 m, empty shells down to 305 m.

Remarks A protoconch of 3.2 conical whorls measured 810 µm high and 700 µm wide at the base. Protoconch-I consisted of 0.9 whorls sculptured with pustules over the entire surface. Protoconch-II consisted of 2.3 whorls with two spiral keels sculptured by a series of subsutural nodulose prosocline threads and a series of opisthocline nodulose threads over each keel. The threads frequently appear to originate from the fusion of two-three tubercles. This protoconch is of the same type as that seen in the Philippines specimen examined. I am unable to separate *B. cookae* from Hawaii (Figs 27, 27) from within the variation of *B. takahashii* and I consider them as synonyms. Furthermore, *B. garrardi* (D'Attilio, 1968) from northern Australia is also similar and it may well be another synonym of *B. takahashii*. The published depth ranges for *B. takahashii* indicating relatively shallow waters (50-120 m) are congruent with the Marquesas records (alive in 90-92 m). The data for Hawaii (136-318 m), as

B. cookae) are more congruent with the dealers' data from the Philippines (alive in 150-300 m). Only by this indication do I classify *B. takahashii* as a deep water species, (Table 1) albeit with caution.

Genus *Leptoconchus* Rüppell, 1834

Leptoconchus Rüppell, 1834: 105. Type species (by subsequent monotypy of Rüppell, 1835): *Leptoconchus striatus* Rüppell, 1835; Recent, Red Sea.

Magilopsis G.B. Sowerby III, 1919: 77. Type species (by original designation): *Leptoconchus lamarckii* Deshayes, 1863; Recent, Indo-Pacific.

Remarks Massin (1982, 1983, 1990, 1992, 2000) studied the ecology and taxonomy of the species in this group. A revision of the *Leptoconchus* species, currently being undertaken by Arjan Gittenberger at the National Museum of Natural History (Naturalis, Leiden, the Netherlands), is revealing a cryptic, adaptive radiation in the genus at least within the species associated with mushroom corals (Fungiidae). Identification of the species seems possible only by molecular markers and/or data on the associated coral (Gittenberger 2006).

Leptoconchus lamarckii Deshayes, 1863

Figures 28, 29

Leptoconchus lamarckii Deshayes, 1863: 127, pl. 13, figs 1, 3.

Type material Five syntypes MNHN.

Type locality Bourbon [Réunion Island].

Material examined MUSORSTOM 9: Ua Pou stn DW 1143, 9°20.9'S, 140°02.7'W, 18-55 m, 3 sh.

Distribution Throughout the Indo-West Pacific (uncommon in the Seychelles: Jarrett 2000), including Hawaii (Kay 1979). New Caledonia and Loyalty Islands, empty shells in 2-60 m.

Remarks Only a few shells of *L. lamarckii* are present in the samples studied, despite it being considered the most common *Leptoconchus*

Table 1 Species of Coralliophilinae from Marquesas Islands, with their record in T&C (Tröndlé & Cosel, 2005) and in the present revision, along with their larval development and depth range (†: based on empty shells only) in the Indo-Pacific (summarized mostly after Oliverio in press) and Marquesas Islands. Habitat: shallow, <100-150 m; deep, >100-150 m (see text for details).

species	T&C	Present	Larval development	Indo-Pacific	Depth range Marquesas	Habitat
<i>Coralliophila violacea</i> (Kiener, 1836)	*	*	P	4-25	0-3 m	shallow
<i>Coralliophila bulbiformis</i> (Conrad, 1837)	*	*	P	4-32	intertidal	shallow
<i>Coralliophila abnormis</i> (Smith, 1878)		*	P	15-25	102-119	shallow
<i>Coralliophila clathrata</i> (A. Adams, 1854)	*	*	P	intertidal-98	intertidal	shallow
<i>Coralliophila monodonta</i> (Blainville, 1832)	*	*	P	intertidal-30	intertidal-33	shallow
<i>Coralliophila curta</i> G.B. Sowerby III, 1894	*		P	intertidal-30	60-62	shallow
<i>Coralliophila nukuhiva</i> n. sp.	*		P?	408-444 †	220-416	deep
<i>Coralliophila mitraiforma</i> Kosuge, 1985	*		P	245-400	1150-1250	deep
<i>Coralliophila nodosa</i> (A. Adams, 1854)	*		P	230-266	104-109	deep
<i>Coralliophila fimbriata</i> (A. Adams, 1854)	*		P	90-305	90-367	deep
<i>Rhizochilus</i> cf. <i>antipathum</i> Steenstrup, 1850	*	*	P	25-400	120-600	deep
<i>Babelomurex japonicus</i> (Dunker, 1882)		*	P	250-253	260	deep
<i>Babelomurex takahashii</i> (Kosuge, 1979)		*	P	50-318 m	90-92	deep?
<i>Leptoconchus lamarcii</i> Deshayes, 1863	*		?	2-60	18-55	shallow

species in the Indo-West Pacific (Massin 1982). More focused sampling on coral boring gastropods would probably give a better idea of their regional diversity.

CONCLUSIONS

Tröndlé & Cosel (2005) reviewed the bibliographic records of molluscs from Marquesas Islands (Hombron & Jacquinot, 1852; Jardin, 1858; Rehder, 1969, 1985; Salvat & Rives, 1975; Tröndlé, 1989), and listed 5 species of Coralliophilinae. Of these, one (*C. bulbiformis*) was not directly collected during the MUSORSTOM expedition, but is reliably recorded from shells in private collections. A total of 14 species have been identified and are treated in the systematic account.

In terms of species richness, the total number of species (14) is in agreement with the total estimate for French Polynesia by Tröndlé (1989), who listed 14 species, with an apparently higher rate of endemism for southern Polynesia. Nevertheless, this survey (part of the MUSORSTOM 9 expedition) used intensive sampling which increased the number of deep water species. The total number of species is about 15% the total number of species from the New Caledonia area (Oliverio in press), and it is 50% of the number of species from southern French Polynesia (Australes Islands: Oliverio in prep.), another marginal area in the Pacific. None of the species is apparently

endemic to Marquesas, and all species probably have a planktotrophic larval development.

The Coralliophilinae display a significant radiation in deep water. Deep habitats in the Indo-West Pacific are considered to be those deeper than 100-150 m, which is the lower limit for the hermatypic scleractinians and where the latter are outcompeted by Alcyonaria, Stylasterids and Porifera. At least seven species can thus be considered as shallow water (including *C. abnormis*, but, conservatively, not *C. fimbriata*). The proportion of deep water species (50%) is significantly smaller than is usual for the Coralliophilinae (65-80%, with an average on a global scale of 75%: Oliverio unpublished). This may be an artefact of the small number of species. Alternatively, it may be an indication that marginality of Marquesas affects the deep water faunas more than the shallow water ones. For epizoic organisms, such as the coralliophilines, this can in turn be related to a reduced availability of suitable hosts in deep waters. The total lack of species with non-planktotrophic development or with a severely shortened pelagic larval phase (which together constitute 10-20% of the Indo-Pacific species: Oliverio in press) seems to support this interpretation. A stronger effect of marginality in deep water is a hypothesis to test with larger databases, when more faunistic inventories are available.

ACKNOWLEDGEMENTS

I wish to thank Philippe Bouchet (MNHN, Paris) who allowed me to revise the coralliophiline material from the French expeditions to Pacific. Financial support of the European Commission's Research Infrastructure Action via the SYNTHESYS Project, is here acknowledged. SEM work was carried out at the "Service commun de Microscopie Electronique des Laboratoires des Sciences de la Vie" (MNHN). The whole LBIMM staff at the "Labò" contributed to make the various stays at the Muséum National d'Histoire Naturelle both proficuous and comfortable. Particularly, Virginie Héros and Philippe Maestrati (MNHN, Paris) are heartily thanked for their assistance with sample management, locality data and locating type material. Jean Tröndlé (La Force) and Michel Boutet (Tahiti) shared information on their collections in French Polynesia. Philippe Bouchet and Pierre Lozouet (MNHN) are thanked for useful discussions on Pacific fauna. Bruce Marshall (Wellington), Ole S. Tendal (ZMUC), Bret Raines (Victorville), Kathie Way and Amelie Machallan (NHM, London), and Thomas von Rintelen (ZMB) are thanked for bibliographic support and type material information and loaning.

REFERENCES

- ADAMS A 1854 Descriptions of new species of *Semele*, *Rhizochilus*, *Plotia* and *Tiara* in the Cumingian Collection. *Proceedings of the Zoological Society of London* **21**[1853]: 94-99.
- ADAMS H & ADAMS A 1853-1858 *The genera of Recent Mollusca; arranged according to their organisation*. 1, 484 pp [1853-54]. 2, 660 pp [1854-58]. 3, plates 1-138 [1858]. J. Van Voorst, London.
- ADAMS H & ADAMS A 1863 Descriptions of new species of shells, chiefly from the Cumingian collection. *Proceedings of the Zoological Society of London* **1863**: 428-435.
- BARNARD KH 1969 Contributions to the knowledge of the South Africa marine Mollusca. Part VI. Supplement. *Annals of South African Museum* **47**: 595-661.
- BENTON TG & SPENCER T 1995 Biogeographic processes at the limits of the Indo-West Pacific Province. In, Benton T.G. & Spencer T. [eds], "The Pitcairn Islands: biogeography, ecology and prehistory". *Biological Journal of the Linnean Society* **56**: 243-244.
- BLAINVILLE H 1832 Disposition methodique des espèces récentes et fossiles des genres Pourpre,
- Ricinule, Licorne et Concholépas de M. de Lamarck. *Nouvelles Annales du Muséum d'Histoire Naturelle Paris* **1**: 189-263.
- BOSCH DT, DANCE SP, MOOLENBEEK RG & OLIVER PG 1995 *Seashells of Eastern Arabia*. Motivate, Dubai.
- BROOK FJ 1999 The coastal scleractinian coral fauna of the Kermadec Islands, southwestern Pacific. *Journal of the Royal Society of New Zealand* **29**: 435-460.
- CERNOHORSKY WO 1972 *Marine shells of the Pacific*. 2. Pacific Publications, Sydney.
- CERNOHORSKY WO 1978 *Marine shells of the Pacific*. 3. Pacific Publications, Sydney.
- CERNOHORSKY WO 1985 The taxonomy of some Indo-Pacific Mollusca. Part 12. With remarks on two American gastropod species. *Records of the Auckland Institute and Museum* **22**: 51-53.
- CHEMNITZ JH 1788, see MARTINI FHW & CHEMNITZ JH 1769-1795.
- CHENU J-C 1859 *Manuel de Conchyliologie et de Paléontologie conchyliologique*. Masson, Paris.
- COEN G 1922 Del genere *Pseudomurex*, (Monterosato 1872). *Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale* **61**: 68-71.
- CONRAD TA 1837 Description of new marine shells from Upper California, collected by Thomas Nuttall, Esq. *Journal of the Academy of Natural Sciences of Philadelphia* **7**: 227-268.
- DALL WH 1924 Notes on Molluscan nomenclature. *Proceedings of the Biological Society of Washington* **37**: 87-90.
- D'ATTILIO A & MYERS BW 1984 Descriptions of five muricacean gastropods and comments on two additional species, in the families Muricidae and Coralliophilidae: (Mollusca). *Transactions of the San Diego Society of Natural History* **20**: 81-94.
- D'ATTILIO A 1968 New species of *Latiaxis* (Gastropoda: Magilidae) from Queensland, Australia. *Journal of the Malacological Society of Australia* **1**(11):7-9. in full please
- DESHAYES GP 1863 Catalogue des Mollusques de l'Ile de Réunion (Bourbon). In: MAILLARD L. (ed.) *Notes sur l'Ile de Réunion*. Annexe E, 6+144 pp, pls 28[1]-41[14]. Dentu, Paris.
- DUNKER W 1882 *Index molluscorum maris Japonici conscriptus et tabulis iconum XVI illustratus*. 301 pp, 16 pls, Theodor Fischer, Cassel.
- FINET Y 1994 *The marine mollusks of the Galapagos Islands: a documented faunal list*. Muséum d'Histoire Naturelle de Genève, Recanati 180 pp.
- GAGE R 1962 New geographical records for Hawaii. *Hawaiian Shell News* **11**(1): 3.
- GITTENBERGER A 2006 *The evolutionary history of parasitic gastropods and their coral hosts in the Indo-Pacific*. PhD Thesis at Leiden University, 249 pp. Leiden
- GMELIN JF 1791 *Caroli Linnaei systema Naturae per regna tria naturae. Editio decima tertia, aucta, reformata, Vermes Testacea*. Leipzig [Lipsiae] **1** (6): 3021-3910 [molluschi].
- HIGO S, CALLOMON P & GOTÖY 1999 *Catalogue and bibliography of the marine shell-bearing mollusca of Japan. Gastropoda, Bivalvia, Polyplacophora, Scaphopoda*. Elle

- Scientific Publications, Ōsaka, Japan, 749 pp.
- HIGO S, CALLOMON P & GOTOY 2001 Catalogue and bibliography of the marine shell-bearing mollusca of Japan. *Gastropoda, Bivalvia, Polyplacophora, Scaphopoda. Type figures*. Elle Scientific Publications, Ōsaka, Japan, 208 pp.
- HOMBON JB & JACQUINOT CH 1852 Zoologie 5. *Voyage au pôle sud et dans l'Océanie sur les corvettes l'Astrolabe et la Zélée, exécuté par ordre du roi pendant les années 1837-1838-1839-1840, sous le commandement de J. Dumont d'Urville, capitaine de vaisseau, publié par ordonnance de Sa majesté*. Pp 132. Gide et Baudry, Paris.
- HUPÉ M 1860 Note sur un genre nouveau de gastéropode, galéropside. *Revue et Magasin de Zoologie* **1860**: 125-128.
- ICZN 1970 Opinion 911. Six misidentified type-species in the superfamily Muricacea (Gastropoda). *Bulletin of Zoological Nomenclature* **27**: 20-26.
- ICZN 1999 *International Code of Zoological Nomenclature. Fourth Edition*. i-xxx + 1-306. ITZN, Padova.
- IREDALE T 1912 New generic names and new species of marine Mollusca. *Proceedings of the Malacological Society of London* **10**: 217-228.
- IREDALE T 1929 Mollusca from the continental shelf of eastern Australia. Part 2. *Records of the Australian Museum* **17**: 157-189.
- JARDIN É [D. Stanislas Aimé] 1858 Essai sur l'histoire naturelle de l'Archipel de Mendana ou des Marquises. 3ème Partie : Zoologie. *Mémoires de la Société Impériale des Sciences Naturelles de Cherbourg* **6**: 161-200.
- JARRETT A 2000 *Marine shells of the Seychelles*. Carole Green Publishing, 149 pp.
- JOUSSEAUME F 1884 Description d'espèces et genres nouveaux de Mollusques. *Bulletin de la Société zoologique de France* **8** [1883]: 186-204, pl 10.
- KAY EA 1965 Marine molluscs in the Cuming collection, British Museum (Natural History) described by William Harper Pease. *Bulletin of the British Museum (Natural History), Zoology, Supplement* **1**: 1-96, pls 1-14.
- KAY EA 1979 *Hawaiian marine shells. Reef and shore fauna of Hawaii, Section 4: Mollusca*. Bernice Pauahi Bishop Museum Special Publication **64(4)**: i-xviii, 1-653.
- KEEN AM 1971 *Sea shells of tropical West America. Marine mollusks from Baja California to Peru*. Second edition. Stanford University Press, Stanford, California, 1064 pp.
- KIENER LC 1835-1836 *Spécies général et iconographie des coquilles vivantes, comprenant la collection du Muséum d'Histoire naturelle de Paris, la collection Lamarck, celle du Prince Massene, (Appartenant maintenant à M. le baron Benjamin Delessert) et les découvertes récentes des voyageurs. 8, Genre Pourpre. (Purpura, Lam.). J.B. Bailliére, Paris.*
- KILBURN RN 1972 Taxonomic notes on South African marine Mollusca (2), with the description of new species and subspecies of *Conus*, *Nassarius*, *Vexillum* and *Demouilia*. *Annals of the Natal Museum* **21**: 391-437.
- KOSUGE S & SUZUKI M 1985 Illustrated catalogue of *Latiaxis* and its related groups. Family Coralliophilidae. *Institute of Malacology of Tokyo, Special Publications* **1**: 1-83.
- KOSUGE S 1979 Descriptions of two new subgenus [sic] and seven new species of the genus *Latiaxis* (Gastropoda, Mollusca). *Bulletin of the Institute of Malacology of Tokyo* **1**: 3-8.
- KOSUGE S 1985 Descriptions of fourteen new species of the family Coralliophilidae from the Western Pacific. *Bulletin of the Institute of Malacology of Tokyo* **2(3)**: 48-57.
- KOSUGE S 1986 Notes on the family Coralliophilidae from South Africa with a description of new species. *Bulletin of the Institute of Malacology of Tokyo* **2 (4)**: 67-69.
- KOSUGE S 1988 Descriptions of two new species of the family Coralliophilidae from Hawaiian Sea (Gastropoda). *Bulletin of the Institute of Malacology of Tokyo* **2 (6)**: 98-100.
- KOSUGE S 1993 Notes on the coralliophilid species from Somalia, East Africa (Gastropoda). *Bulletin of the Institute of Malacology of Tokyo* **3 (1)**: 9.
- KURODA T 1931 New Japanese shells. Part 3. *Venus* **2**: 314-318.
- KURODA T 1953 New genera and species of Japanese Rapidae. *Venus* **17**: 117-130.
- LAMARCK JBPA DE M DE 1816 Liste des objets représentés dans les planches de cette livraison. «*Vingt-troisième partie: mollusques et polypes divers, par M. Lamarck. In Bruguiere JG, Lamarck JBP & Vincent B de St (1782-1832) Tableaux encyclopédique et méthodique des trois règnes de la nature*». 1-16, pls 391-488, 431 bis, 432 bis*.
- LAN TC 1982 A new coralliophilid species from Bohol Island, Philippines. *Bulletin of Malacology Republic of China* **9**: 23-25.
- LOZOUET P & RENARD P 1998 Les Coralliophilidae, Gastropoda de l'Oligocène et du Miocène inférieur d'Aquitaine (Sud-Ouest de la France): systématique et coraux hôtes. *Geobios* **31**: 171-184.
- MARTINI FHW & CHEMNITZ JH 1769-1795 *Neues Systematisches Conchylien-Cabinet, geordnet und beschreiben von Friedrich Heinrich Wilhelm Martini [vols. 1-3], ... forgesetz und nun vollendet von Johann Heironymous Chemnitz [vols. 4-11] ... Vol. 1: i-xxviii + 1-408, vigs 1-16, pls 1-31 [1769]; Vol. 2: i-xvi + 1-362, vigs 17-26, pls 32-65 [1773]; Vol. 3: i-vi + 1-434, vigs 27-33, 33bis, 34-36, 36bis, 66-121 [1777]; Vol. 4: , vigs 37-41, pls 122-159 [1780]; Vol. 5: 1-324, vigs 42-44, pls 160-193 [1781]; Vol. 6: 1-375, vigs 1-7, pls 1-36 [1782]; Vol. 7: 1-356, vigs 8-11, pls 37-69 [1784]; Vol. 8: 1-372, vigs 12-18, pls 70-102 [1785]; Vol. 9: vig. 19, pls 103-136 [1786]; Vol. 10: 1-310, vigs 20-25, pls 137-173 [1788]; Vol. 11: 1-310, vig. 26, pls 174-213 [1795]. Gabriel Nikolaus Raspe, Nürnberg, Germany.*
- MASSIN C 1982 Contribution to the knowledge of two boring gastropods with an annotated list of the genera *Magilus* Montfort, 1810 and *Leptoconchus* Rüppell, 1835. *Bulletin de l'Institut Royal de Sciences Naturelles de Belgique, Biologie* **53**: 1-28 + 1 pl.

- MASSIN C 1983 Notes on the genus *Leptoconchus* Rüppell, 1835 (Mollusca, Gastropoda, Coralliophilidae) with the description of two new species, *Leptoconchus vangoethemi* sp.n. and *Leptoconchus cyphastreæ* sp.n., from Papua New Guinea. *Bulletin de l'Institut royal de Sciences naturelles de Belgique, Biologie* **55**: 1-16 + 5 pls.
- MASSIN C 1987 *Reliquiaecava*, a new genus of Coralliophilidae (Mollusca, Gastropoda). *Bulletin de l'Institut Royal de Sciences Naturelles de Belgique, Biologie* **57**: 79-90.
- MASSIN C 1990 Biologie ed écologie de *Leptoconchus peronii* (Lamarck, 1818) (Gastropoda, Coralliophilidae) récolté en Papouasie Nouvelle-Guinée, avec une redescription de l'espèce. *Bulletin de l'Institut royal de Sciences naturelles de Belgique, Biologie* **60**: 23-33.
- MASSIN C 1992 Quelques aspects de la biologie des Coralliophilidae et plus particulierement du genre *Leptoconchus*. *Apex*, hors série: 18-20.
- MASSIN C 2000 Ecology of *Leptoconchus* spp. (Gastropoda, Coralliophilidae) infesting Fungiidae (Anthozoa, Madreporearia). *Bulletin de l'Institut royal de Sciences naturelles de Belgique, Biologie* **70**: 235-252.
- MONTEROSATO TA DE M, DI 1872 *Notizie intorno alle conchiglie fossili di Monte Pellegrino e Ficarazzi*. 44 pp. Michele Amenta, Palermo.
- OLIVER WRB 1915 The Mollusca of the Kermadec Islands. *Transactions of the New Zealand Institute* **47**: 509-568.
- OLIVERIO M in press Coralliophilinae (Neogastropoda: Muricidae) from the south-west Pacific. Tropical Deep Sea Benthos (in press).
- OLIVERIO M & MARIOTTINI P 2001 A molecular framework for the phylogeny of Coralliophila and related muricoids. *Journal of Molluscan Studies* **67**: 215-224.
- OLIVERIO M, CERVELLI M & MARIOTTINI P 2002 ITS2 rRNA evolution and its congruence with the phylogeny of muricid neogastropods (Caenogastropoda, Muricoidea). *Molecular Phylogenetics and Evolution* **25**: 63-69.
- PEASE WH 1861 Descriptions of seventeen new species of marine shells, from the Sandwich islands, in the collection of H. Cuming. *Proceedings of the zoological Society of London* **1860**: 397-400.
- PEASE WH 1865 Descriptions of new genera and species of marine shells from the islands of the Central Pacific. *Proceedings of the zoological Society of London* **1865**: 512-517.
- QUOY JCR & GAIMARD P 1832-1835 *Voyage de découvertes de l'Astrolabe, exécuté par ordre du Roi pendant les années 1826-1827-1828-1829, sous le commandement de M. J. Dumont d'Urville*. Zoologie. 2[1832]: 1-320; 2[1833]: 321-686; 3[1834]: 1-366; 3[1835]: 369-954, 8 pls [including the "Table générale des planches"]. Tastu, Paris.
- RAFINESQUE CS 1815 *Analyse de la nature ou tabluae de l'univers et des corps organisée*. Palermo. 223 pp.
- REEVE LA 1846 Genus *Purpura*. *Conchologia Iconica: figures and descriptions of the shells of molluscs; with remarks on their affinities, synonymy, and geographical distribution*. 3, Reeve, Brook, London.
- REHDER HA 1969 The marine molluscan fauna of the Marquesas Islands. *Annual Report for 1968 of the American malacological Union* **1968**: 29-32.
- REHDER HA 1980 The marine mollusks of Easter Island (Isla de Pascua) and Sala y Gómez. *Smithsonian Contributions to Zoology* **289**: 167 pp.
- REHDER HA 1985 A new species of Coralliophila (Gastropoda: Coralliophilidae) from Southeastern Polynesia. *The Nautilus* **99**: 97-100.
- RICHER DE FORGES B, POUPIN J & LABOUTE P 1999 La campagne MUSORSTOM 9 dans l'archipel des île Marquises (Polynésie française). Compt rendu et liste des stations. In Crosnier A. (ed.) *Résultat des Campagnes MUSORSTOM*, Volume 20. *Mémoires du Muséum National d'Histoire Naturelle* **180**: 9-29.
- ROBERTSON R 1980 *Epitonium millecostatum* and *Coralliophila clathrata*: two prosobranch gastropods symbiotic with Indo-Pacific *Palythoa* (Coelenterata: Zoanthidae). *Pacific Science* **34**: 1-17.
- RÜPPELL E 1834 Description of a new genus of pectinibranchiated gasteropodous Mollusca (*Leptoconchus*). *Proceedings of the Zoological Society of London* **1834**: 105-106.
- RÜPPELL E 1835 Description d'un nouveau genre de mollusques de la classe des gastéropodes pectinibranches. *Transactions of the Zoological Society of London* **1**: 259-260.
- SALVAT BG & RIVES C 1975 *Coquillages de Polynésie*. Les Editions du Pacifique. Papeete, Tahiti. 391pp.
- SHIKAMA T 1963 On some noteworthy marine Gastropoda from Southwestern Japan. *Science Reports of the Yokohama National University* sect. 2, 10: 61-66, 1 fig., pl. 1.
- SMITH EA 1878 On a collection of marine shells from the Andaman Islands. *Proceedings of the Zoological Society of London* **1878**: 804-821, pl. 50.
- SOUVERBIE M 1861 Descriptions d'espèces nouvelles de l'archipel Calédonien. *Journal de Conchyliologie* **9**: 271-284.
- SOWERBY GB III 1894 Descriptions of twelve new species, chiefly from Mauritius. *Proceedings of the Malacological Society of London* **1**: 41-44, pl. 4.
- SOWERBY GB III 1919 Notes on *Magilus* and its allies, substituting the generic name *Magilopsis* for *Leptoconchus lamarcki*, Deshayes. *Proceedings of the Malacological Society of London* **13**: 75-77.
- SOWERBY J & SOWERBY GB I 1821-1834 *The genera of Recent and fossil shells, for the use of students in conchology and geology*. Vol. 1: pls 1-126 + unnumbered pages [1821-25]; vol. 2: pls 127-262 + unnumbered pages [1825-34], London.
- STEENSTRUP JJS 1850 Beskrivelse over nogle maerkelige tildeelde nye Dyrarter. *Oversigt over det Kongelige Danske Videnskabernes Selskabs Forhandlinger* **1850**: 75-77.
- TRÖNDLÉ J & VON COSEL R 2005 Inventaire bibliographique des mollusques marins de l'Archipel des Marquises (Polynésie française). *Atoll Research Bulletin* **542**: 265-339.

- TRÖNDLÉ J 1989 Coralliophilidae de Polynésie Française. *Xenophora* **45**: 18-27.
- TSUCHIYA K 2000 Superfamily Muricoidea. Pp 365-422. In: OKUTANI, T. (ed.) *Marine mollusks in Japan*. Tokai University Press, Tokyo.
- WOOLACOT L 1954 New shells from New South Wales. *Proceedings of the Royal Society of New South Wales* **1952-53**: 37-39.
- YEN TC 1942 A review of Chinese gastropods in the British Museum. *Proceedings of the Malacological Society of London* **24**: 170-289, pls 11-28.