ASHFORDIA GRANULATA (HYGROMIIDAE) LIVING IN NORTHERN FRANCE

Ashfordia granulata (Alder, 1830) is primarily known from Great-Britain¹, with a few records from Brittany², Cotentin³ and also from N.W. Spain⁴. According to Kerney and Cameron², the distribution of this species is lusitanic.

In February 2004 a small, but stable, population of Ashfordia granulata was recorded from Fort-Mahon-Plage (Dept. Somme), 30 km West from Abbeville. All the specimens were observed

in the leaf litter of the Common Reed (Phragmites australis), at the foot of a sea wall defence. All living specimens observed were juvenile snails and all the adults were dead shells. Amongst other common associates of Ashfordia granulata a stablepopulation of Vertigo angustior was noted at this new site. The habitat was similar to those ofpopulations found in eastern Great Britain. According to Kerney¹, the Silky snail is restricted

in eastern Great Britain to marshy ground in river valleys and this recent find is located in the Bay of Authie River, a small coastal river.

This locality thus represents a considerable extension of the continental distribution of the species to the North and provides the most northern population in France to date.

¹Kerney, M.P. 1999 Atlas of the land and freshwater Molluscs of Britain and Ireland

²Kerney & Cameron 1999 *Guide des escargots et limaces d'Europe* Delachaux & Niestlé, Lausanne.

³MAZURIER, M. & COCHARD, P.-O 2003 Lettre de Germain 4: 1-6.

⁴ALTONOGA, K. et al. 1994 Estudio faunistico y biogeographico de los Moluscos terrestres del norte de la Peninsula Iberica.

Xavier Cucherat
GREET Ing
Route du Musée, Haringzelles
F-62179 Audinghen
E-mail: xavier.cucherat@wanadoo.fr
François Boca
Conservatoire des Sites Naturels de Picardie
1, place Ginkgo - village Oasis
F-80044 Amiens Cedex 1
E-mail: f.boca@conservatoirepicardie.org